



CYTHRICIN 25%

CYPERMETHRIN - HIGH CIS 25% EMULSIFIABLE



COMPOSITION :

Each ml contains :

Cypermethrin - High Cis 25% w/v
Solvents and emulsifiers q.s.

INDICATION :

Acts on wide range of Ectoparasites such as Flies, Lice, Mites, Ticks and other Ectoparasites on Cattle, Pigs, Poultry, Sheep and Goats.

DOSAGE AND ADMINISTRATION :

For external use only.

Dilute the CYTHRICIN 25 % water using 150 times the volume of water, mix well until a milky emulsion is produced.

Prepare fresh solution every time. Product is suitable for application with all types of surface spraying equipment.

For Cattle :

For Tick control : All stages of single and multi-host ticks : Mix 3 ml of CYTHRICIN 25% with 5 litres of water. Apply to the animal until thoroughly wet specific to areas where ticks are attached. Check application at specific areas where ticks are attached. Spraying should require 3-5 times per animal.

For Mite control : Mix 2 ml CYTHRICIN 25% 5 litres of water. Spray 250 ml mixture to each side of Adult Cattle and 75 ml per side of Calves. On fly season repeat every 28 days for controlling of biting flies and at 14 days for fly control. CYTHRICIN 25% is safe to use on lactating dairy animals without the need to discard milk from treated animals. Care should be taken to prevent direct contamination of milk or equipment with spray mixture.

For Sheep :

For Mites and Lice control : Mix 2 ml of CYTHRICIN 25% with 10 litres water and dip or spray the animal. For heavy infestations of mites increase the strength of mixture to 4 ml per 10 litres.

For Goats :

For Ticks, Mange, Lice, Nuisance flies control : Mix 3 ml CYTHRICIN 25% with 10 litres of water and spray to thoroughly wet the animal.

For Pigs :

For Ticks, Mange, Lice, Nuisance flies control : Mix 3 ml CYTHRICIN 25% with 10 litres of water and spray till animal gets wet.

For Poultry :

For Lice and Mites control : Mix 2 ml of CYTHRICIN 25% with 1 litre of water and apply directly to birds and to all surfaces of houses. Repeat every 28 days as necessary.

Do not apply to newly hatched chicks.

For Animal Houses : For control of nuisance flies : Mix 2 ml CYTHRICIN 25% with 1 litre of water and spray at a rate of 1 litre per 10 squaremetres. Suitable in Poultry houses, farrowing houses, dairy parlours and all farm buildings.

Ectoparasites



WITHDRAWAL PERIOD :

Cattle : Meat - 28 days, Milk - Nil
Sheep and Goats : Meat - 18 days
Poultry : Meat - 21 days : Egg - Nil

STORAGE :

Store below 30°C in a dry place, protect from direct sunlight.

PACK SIZE :

1 litre.

Manufactured in India by :

IPPL
Intracin
Pharmaceuticals Pvt. Ltd.
C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.



Ectoparasites