



DONECIN



DOXYCYCLINE AND NEOMYCIN POWDER FOR ORAL ADMINISTRATION

COMPOSITION :

Each gm contains :

Doxycycline hyclate	B.P.	
Equivalent to Doxycycline		100 mg
Neomycin sulphate	B.P.	
Equivalent to Neomycin		100 mg
Excipients		q.s.

INDICATIONS :

Poultry : For treatment of mixed infections specially complicated CRD, Nonspecific diarrhoea, Blue comb, Mud fever, Fowl cholera, Coli-septicemia, Pullorum disease, infectious synovitis. Secondary bacterial infections caused by gram positive and gram negative bacteria.

DOSAGE AND ADMINISTRATION :

Poultry : Treatment of severe outbreaks :

1 gm in 5 litres of drinking water for 4-5 days or till infection is eliminated.

CONTRAINDICATIONS :

Hypersensitivity to Tetracyclines and/or Neomycin.

Concurrent administration of Penicillinase, Cephalosporins, Quinolones, Cycloserine and Bacteriostatic agents.

Overdosage, due to small safety margin of Neomycin.

Do not use in combination with Cephalosporin or Penicillin due to increased risk of renal damage.

WITHDRAWAL PERIOD :

The withdrawal period after the last dose/treatment

is : Poultry : Meat : 7 days

PACK SIZE :

50 g, 100 g & 250 g Sachet.

Manufactured in India by :



Intracin
Pharmaceuticals Pvt. Ltd.

C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

