

FLASIN 20%

TYLOSIN INJECTION B.P. 20% W/V



COMPOSITION :

Each ml contains :

Tylosin Tartrate	B.P.	
equivalent to Tylosin base		200 mg
Benzyl alcohol	B.P.	2% v/v
Propylene glycol	B.P. and	
Water for Injection	B.P. (1:1)	q.s.

INDICATIONS :

Cattle : Tylosin Injection is indicated for use in the treatment of bovine respiratory complex (shipping fever, pneumonia) usually associated with *Pasteurella multocida* and *Actinomyces pyogenes*; foot rot (necrotic pododermatitis) and calf diphtheria caused by *Fusobacterium necrophorum* and metritis caused by *Actinomyces pyogenes*.

Sheep : Pneumonia, arthritis, contagious and acute agalactia due *Mycoplasma agalactiae*, mastitis.

Goats : Contagious pleuropneumonia due *Mycoplasma mycoides*.

CONTRAINDICATION :

Not to be used in animals known to be hypersensitive to the active ingredient.

DOSAGE AND ADMINISTRATION :

Cattle :

Inject 5-10 mg/kg (0.025-0.05 ml/kg) of bodyweight daily Intramuscularly.

Do not Inject more than 10 ml per Injection site.

Sheep and Goats :

Inject 10 mg/kg (0.05 ml/kg) of bodyweight daily Intramuscularly.

Do not Inject more than 5 ml per Injection site.

WITHDRAWAL PERIOD :

Cattle :

Edible tissue : 7 days

Milk : 72 hours

Sheep, Goats :

Edible tissue : 7 days

Milk : 48 hours

PACK SIZE : 30 ml & 100 ml Vials.

Manufactured in India by :

IPPL
Intracin
Pharmaceuticals Pvt. Ltd.
C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

