



FLASIN 20% WS

TYLOSIN POWDER FOR ORAL ADMINISTRATION



COMPOSITION :

Each gm contains :

Tylosin tartrate B.P. (vet.) equivalent to

Tylosin 20% w/w

Excipients q.s.

Each 100 g sachet and 500 gm & 1 kg Jar of FLASIN 20% WS contains 20 g Tylosin.

INDICATIONS :

FLASIN 20 % WS is indicated in all conditions associated with bacteria sensitive to Tylosin.

This includes organisms in the following genera :

Streptococcus, Bacillus, Staphylococcus, Corynebacterium, Clostridium, Erysipelothrix, Spirochaetes, Fusiformis, Chlamydia, Mycoplasma, Pasteurella and Campylobacter.

CONTRAINDICATIONS :

Not to be used in animals hypersensitive to Tylosin or other macrolide antibiotics e.g. Erythromycin.

Do not use FLASIN 20 % WS in layers producing eggs for human consumption.

DOSAGE AND ADMINISTRATION :

Pigs : 3-6 mg Tylosin (0.015 g to 0.03 g of FLASIN 20 % WS) per kg bodyweight.

Calves : 0.5 g to 1 g Tylosin (2.5 g to 5 g of FLASIN 20 % WS) per Calf administered orally twice daily for 7-14 days. The product may be given in the milk or milk replacer.

Chickens and Turkeys : 0.5 g Tylosin (2.5 g of FLASIN 20 % WS) in 1 litre of drinking water.

WITHDRAWAL PERIOD :

Animals must not be slaughtered for human consumption during treatment.

The withdrawal periods after the last treatment are as follows :

Meat :

Calves : 14 days

Broiler chickens : 1 day

Turkeys and Pigs : Nil

PACK SIZE :

100 gram Sachet and 500 gram & 1 kg Jar.

Manufactured in India by :



Intracin

Pharmaceuticals Pvt. Ltd.

C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

