



FLASIN WS

TYLOSIN TARTRATE POWDER FOR ORAL ADMINISTRATION



COMPOSITION :

Each gram contains :

Tylosin tartrate B.P. 1000 mg

INDICATIONS :

Gastrointestinal and respiratory infections caused by Tylosin sensitive micro-organisms like *Campylobacter*, *Mycoplasma*, *Pasteurella*, *Staphylococcus*, *Streptococcus* and *Treponema* spp. in Calves, Poultry, Kids and Lambs.

DOSAGE AND ADMINISTRATION :

For oral administration :

Calves, Kids and Lambs :

Twice daily, 5 g per 220 – 250 kg bodyweight for 5 – 7 days.

Poultry :

1 kg per 1500 – 2000 liters of drinking water for 3-5 days.

Note : for pre-ruminant Calves, Lambs and Kids only.

CONTRAINDICATIONS :

Hypersensitivity to Tylosin.

Concurrent administration of Penicillins, Cephalosporins, Quinolones and Cycloserine.

Administration to animals with an active microbial digestion.

WITHDRAWAL PERIOD :

For Meat :

Calves, Poultry, Kids and Lambs : 5 days

PACK SIZE :

30 g, 100 gram Sachet, 1 kg of Jar.

Manufactured in India by :



Intracin
Pharmaceuticals Pvt. Ltd.

C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

