



I-TIN

IVERMECTIN INJECTION 1% B.P.



COMPOSITION :

Each ml contains :

Ivermectin	B.P.	10 mg
Benzyl alcohol	B.P.	1.5% v/v
(As Preservative)		
Propylene glycol	B.P.	q.s.

INDICATION : For treatment and control of gastrointestinal nematodes, Lung worms, Eye worms, Warbles and Ectoparasites like mites, ticks, and lice in Cattle, Buffaloes, Horses, Sheep, Goats, Camels and Dogs.

CONTRA INDICATION :

I-TIN should not be used Intravenously or Intramuscularly.

Do not use in lactating Cows or in dairy Cows during 28 days prior to calving.

DOSAGE AND ADMINISTRATION :

Cattle, Buffaloes, Horses, Sheep, Goats, Camels and Dogs :

Dosage : 1 ml per 50 kg body weight (based on a recommended level of 200 mcg Ivermectin per kg bodyweight).

Administration : Inject Subcutaneously in front or behind the shoulder.

WITHDRAWAL PERIOD :

Cattle, Buffaloes : Must not be treated within 21 days of slaughter for human consumption. Do not use in Cattle producing milk for human consumption or in dairy Cows within 28 days prior to calving.

Sheep : Must not be treated within 21 days of slaughter for human consumption. Do not treat lactating Cows or within 21 days before lambing where milk is to be used for human consumption.

Horses : Withholding periods before slaughter 35 days.

Goats : Meat : 35 days, Milk : 40 days.

Camels : Must not be slaughtered for human consumption within 28 days from the last day of treatment.

Camels producing milk for human consumption should not be treated.

PACK SIZE :

10 ml, 20 ml, 50 ml, 100 ml, 250 ml Vials.

Manufactured in India by :

IPPL
Intracin
Pharmaceuticals Pvt. Ltd.
C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

