



IMOLY 10%



OXYTETRACYCLINE INJECTION 10% W/V

COMPOSITION :

Each ml contains :

Oxytetracycline dihydrate	B.P. eq. to	
Anhydrous oxytetracycline		100 mg
Propylene glycol	B.P.	q.s.

INDICATIONS :

Arthritis, gastrointestinal and respiratory infections caused by Oxytetracycline sensitive micro-organisms, like Bordetella, Campylobacter, Chlamydia, E.coli, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in Cattle, Calves, Goats and Sheep.

DOSAGE AND ADMINISTRATION :

For Intramuscular or Subcutaneous administration :

Full-grown animals : 1 ml per 10 - 20 kg bodyweight, for 3 - 5 days.

Young animals : 2 ml per 10 - 20 kg bodyweight, for 3 - 5 days.

Do not administer more than 20 ml in Cattle, more than 5 ml in Calves, Goats and Sheep per injection site.

CONTRAINDICATIONS :

Hypersensitivity to Tetracyclines.

Administration to animals with a seriously impaired renal and/or hepatic function.

Concurrent administration of Penicillins, Cephalosporins, Quinolones and Cycloserine.

WITHDRAWAL PERIOD :

For meat : 12 days

For milk : 5 days

PACK SIZE : 50 ml and 100 ml Vials.

Manufactured in India by :

IPPL
Intracin
Pharmaceuticals Pvt. Ltd.
C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

