



IMOLY 20% WS

OXYTETRACYCLINE HYDROCHLORIDE
POWDER 20% W/W



COMPOSITION :

Each gram contains :

Oxytetracycline hydrochloride B.P. 200 mg

INDICATIONS :

IMOLY 20 % powder is indicated for the treatment and control of bacterial infections such as infectious coryza, fowl cholera, infectious synovitis, fowl typhoid, chronic respiratory disease, bacillary white diarrhoea, spirochaetosis and to check secondary bacterial infection during viral outbreaks.

IMOLY 20 % soluble powder is also can be used to treat infectious diseases caused by Pigs and Chickens that are sensitive to Escherichia coli, Salmonella, Pasteurella and Mycoplasma.

1. Diarrhoea caused by Escherichia coli or Salmonella, yellow scour of Piglets, white scour of Piglets, white scour of young Chickens and Calves and Lamb dysentery.
2. Swine asthma, bovine pneumonia and chronic respiratory disease in Chickens caused by Mycoplasma.
3. Swine pneumonia, gallbladder disorder and bovine hemorrhagic sepsis caused by Pasteurella multocida.
4. Fever, pale skin and conjunctiva caused by porcine eperythrocytosis, yellow staining in severe cases, constipation or diarrhoea, general fatigue, lethargy and swollen lymph nodes.

DOSAGE AND ADMINISTRATION :

Chickens & Turkey :

Infectious synovitis caused by Mycoplasma synoviae, susceptible to IMOLY 20 % powder 1-2 g per 4.5 liter of water for 3-5 days.

Chronic respiratory disease and air sac infections caused by Mycoplasma gallisepticum, Salmonellosis and fowl cholera caused by Pasteurella multocida : IMOLY 20 % powder 2-4 g per 4.5 liter of water for 3-5 days.

Calves : 10-20 g per 200 kg body weight every 12 hours during 3-5 days.

Pigs : 1-2 g per liter drinking water during 3-5 days.

WITHDRAWAL PERIOD :

Meat : Calves : 8 days, Pigs : 5 days, Poultry : 8 days.

Eggs : 4 days

PACK SIZE :

100 gm Sachet, 500 gram and 1 kg Jar.

Manufactured in India by :



Intracin

Pharmaceuticals Pvt. Ltd.

C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

