



INCEFT

CEFTIOFUR SODIUM POWDER FOR INJECTION 1G/4G (VET)



COMPOSITION :

Each vial contains :

Ceftiofur sodium sterile powder USP
equi. to Ceftiofur 1000 mg

This pack also contains 20 ml sterile water for Injection I.P. for reconstitution along with sterile syringe and needle.

INDICATION :

Genital infections of bovine (acute metritis, cervicitis, prolapse related to retention of placenta cases) associated with *Trueperella pyogenes*, *Fusobacterium necrophorum* and *Bacteroides* spp. Respiratory diseases of Cattle, Buffaloes, Sheep and Goats (shipping fever, pneumonia) associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni*.

Acute interdigital necrobacillosis (foot rot, pododermatitis) caused by *Fusobacterium* and *Bacteroides*.

DOSAGE AND ADMINISTRATION :

I.M / S.C USE ONLY

Cattle, Sheep & Goats : 1.1 to 2.2 mg / kg b.wt. or 1 to 2 ml (after reconstitution) / 45.45 kg b.wt. by IM use only, once daily for 3-5 days.

Buffaloes : 2 to 2.4 mg / kg b.wt. by IM use only, once daily for 3-5 days.

Dogs : 2.2 mg / kg b.wt. by SC use only, once daily for 5-14 days.

Camels : 2.2 mg / kg b.wt. by IM, once daily for 3-5 days.

Horses : 2.2 to 4.4 mg / kg b.wt. or 1 to 2 ml (after reconstitution) / 22.72 kg b.wt. by IM use only, once daily for 3-5 days.

Swine : 3 to 5 mg / kg b.wt. or 1 to 1.7 ml (after reconstitution) / 16.66 kg b.wt. by IM use only, once daily for 3-5 days.

Chicken (day-old chick) : 0.08 to 0.2 mg / Chick, SC use only in the neck region (1g vial after reconstitution with 20 ml water for injection will treat approximately 5,000 to 12,500 day - old chicks).

WITHDRAWAL PERIOD :

For Cattle, Buffaloes and Swine : Milk : Nil, Meat : 4 days

For Sheep and Goats : Milk : Nil, Meat : Nil

PACK SIZE :

1 gram in 30 ml Vial.

Manufactured in India by :

IPPL
Intracin
Pharmaceuticals Pvt. Ltd.
C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.



Veterinary Medicines - Injectables