



INOXY 20% WS

OXYTETRACYCLINE HCL POWDER

COMPOSITION :

Each gram contains :

Oxytetracycline hydrochloride B.P. 200 mg
Excipients ad 1 gm

INDICATIONS :

INOXY 20% powder is indicated for the treatment and control of bacterial infections such as infectious coryza, fowl cholera, infectious synovitis, fowl typhoid, chronic respiratory disease (CRD), bacillary white diarrhea spirochaetosis and to check secondary bacterial infection during viral outbreaks.

DOSAGE AND ADMINISTRATION :

CHICKENS AND TURKEYS :

Infectious synovitis caused by *Mycoplasma synoviae*, susceptible to Oxytetracycline : 200-400 mg per 4.5 liter of drinking water or INOXY 20% powder 1-2 g per 4.5 liter of water for 3-5 days.

Chronic respiratory disease (CRD) and air sac infections caused by *Mycoplasma gallisepticum*, *Salmonellosis* and fowl cholera caused by *Pasteurella multocida* : 400-800 mg per 4.5 liter of drinking water or INOXY 20% powder 2-4 g per 4.5 liter of water for 3-5 days.

WITHDRAWAL PERIOD :

Meat : 7 days from the last treatment.

PACK SIZE :

100 gram, 500 gram and 1 kg Jar.

Manufactured in India by :

IPPL
Intracrin
Pharmaceuticals Pvt. Ltd.
C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.



Veterinary Medicines - Water Soluble Powders