



IROFLOX



ENROFLOXACIN INJECTION 10% W/V

COMPOSITION :

Each ml contains :

Enrofloxacin	USP	100 mg
Benzyl alcohol	B.P.	1.5% v/v
(As preservative)		
Water for Injection	B.P.	q.s.

INDICATIONS :

Shipping fever pneumonia (H.S.), Pneumonia, Pleuropneumonia, Pseudo rinderpest and Urinary infections, mastitis caused by Mycoplasma spp. Respiratory and enteric diseases (Pasteurellosis, Mycoplasmosis, Colibacillosis, Colisepticaemia and secondary bacterial complication during viral infections). Mastitis caused by susceptible microorganisms and renal infection. IROFLOX is effective against the following microorganisms: Mycoplasma spp., E.coli, Salmonella spp., Haemophilus spp., Pasteurella spp., Klebsiella spp., Proteus spp., Vibrio parahaemolyticus and Staphylococcus aureus.

CONTRAINDICATIONS :

Do not exceed the recommended dose. In accidental overdosage there is no antidote and treatment should be symptomatic.

DOSAGE AND ADMINISTRATION :

As directed by the Veterinarian or 1 ml per 20-40 kg body weight depending upon the severity of infection. (2.5-5 mg of Enrofloxacin 1 kg body weight) Give for 3-5 days.

ROUTE : For IM / IV / SC use.

WITHDRAWAL PERIOD :

Animals must not be slaughtered and milk should not be use for human consumption during treatment. The withdrawal periods after the last treatment are :

Cattle, Camels, Sheep, Goats, Buffaloes :

For Intravenous route : Meat : 4 days, Milk : 72 hours

For Subcutaneous route : Meat : 14 days, Milk : 84 hours

PACK SIZE :

50 ml and 100 ml Vials.

Manufactured in India by :

IPPL
Intracin
Pharmaceuticals Pvt. Ltd.
C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

