

IROFLOX SA



ENROFLOXACIN INJECTION 10% W/V (VET.)

COMPOSITION :

Each ml contains :

Enrofloxacin	USP	100 mg
Benzyl alcohol	B.P.	20 mg
(As preservative)		
Water for Injection	B.P.	q.s.

INDICATIONS :

Shipping fever pneumonia (H.S), Pneumonia, Pleuropneumonia, Pseudo rinderpest and Urinary infections, mastitis caused by Mycoplasma spp. Respiratory and enteric diseases (Pasteurellosis, Mycoplasmosis, Colibacillosis, Colisepticaemia and secondary bacterial complication during viral infections). Mastitis caused by susceptible microorganisms and renal infection. IROFLOX SA is effective against the following microorganisms: Mycoplasma spp., E.coli, Salmonella spp., Haemophilus spp., Pasteurella spp., Klebsiella spp., Proteus spp., Vibrio parahaemolyticus and Staphylococcus aureus.

DOSAGE AND ADMINISTRATION :

Cattle, Buffaloes, Sheep, Goats and Camels :

For Subcutaneous or Intravenous injection.

As directed by the veterinarian or 1 ml per 20-40 kg body weight depending upon the severity of infection. (2.5-5 mg of Enrofloxacin per 1 kg body weight) the second administration should be administered after 72 hours, if required.

WITHDRAWAL PERIOD :

Animals must not be slaughtered and milk should not be used for human consumption during treatment.

Withdrawal periods after the last treatment are :

Cattle :

For intravenous route : Meat : 4 days, Milk : 72 hours (i.e. 6 milkings from the last treatment)

For Subcutaneous route : Meat : 14 days, Milk : 84 hours (i.e. 7 milkings from the last treatment.)

PACK SIZE :

50 ml, 100 ml Vial.

Manufactured in India by :



Intracin

Pharmaceuticals Pvt. Ltd.

C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

