



ISOMID



ISOMETAMIDIUM CHLORIDE HYDROCHLORIDE

COMPOSITION :

Each sachet contains :

Isometamidium chloride hydrochloride 1 g

INDICATIONS :

ISOMID is indicated for treatment and prevention of trypanosomiasis caused by *Trypanosoma* spp. in Cattle, Goats, Sheep, Camels, Horses and Dogs.

A preventive dosage ensures protection for 2 to 4 months.

When clinical cases occur, the whole group should be treated.

DOSAGE AND ADMINISTRATION :

Dissolve the sachet content in to 100 ml of sterile water and administer 1 ml of the solution for 10 kg liveweight (0.5 mg/kg) by deep I.M. or I.V. Injection.

WITHDRAWAL PERIOD :

Meat : 1 month

Milk : Nil

PACK SIZE :

1 gram of Sachet.

125 mg of Sachet.

One sachet is sufficient for the preventive treatment of 8 animals or the curative treatment of 16 animals, weighting 250 kg each.

Miscellaneous - Trypanosomiasis

Manufactured in India by :



Intracin

Pharmaceuticals Pvt. Ltd.

C-1, B-52,53,56, G.I.D.C. ESTATE,

NADIAD - 387 001, (GUJARAT) INDIA.

