



MARBACIN 10%

MARBAFLOXACIN INJECTION

10% W/V B.P.



COMPOSITION :

Each ml contains :

Marbofloxacin B.P. 100 mg

m-cresol B.P. 2 mg

(As preservative)

Water for Injection B.P. q.s.

INDICATIONS :

Cattle : Treatment of respiratory infection caused by sensitive strains of *Pasteurella multocida*, *Mannheimia (Pasteurella) haemolytica* and *Mycoplasma bovis*. It is also indicated in the treatment of acute mastitis caused by *E.coli* strains sensitive to Marbofloxacin during the lactation period.

DOSAGE AND ADMINISTRATION :

Cattle :

Treatment of respiratory Infections :

The recommended dosage is 1 ml/50 kg bodyweight in a single daily injection by Subcutaneous or Intramuscular route for 3 to 5 days.

Treatment of acute mastitis :

The recommended dosage is 1 ml/50 kg bodyweight in a single daily injection by Subcutaneous or Intravenous route for 3 consecutive days.

The first Injection may also be given by the Intravenous route.

CONTRAINDICATIONS :

Bacterial infections with resistance to other fluoroquinolones (cross resistance).

Do not administer to an animal previously found to be hypersensitive to Marbofloxacin or other quinolone.

WITHDRAWAL PERIOD :

Cattle :

Meat and offals : 6 days

Milk : 36 hours

PACK SIZE :

30 ml, 50 ml, 100 ml Vial.

Manufactured in India by :



Intracin

Pharmaceuticals Pvt. Ltd.

C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

