



MELOXIN-I

MELOXICAM INJECTION B.P. (VET) 5MG/ML



COMPOSITION :

Each ml contains :

Meloxicam	B.P.	5 mg
Benzyl alcohol	B.P.	2% v/v
Water for Injection	B.P.	q.s.

INDICATIONS :

Dogs : MELOXIN - I Injection is indicated in Dogs for the control of pain and inflammation associated with osteoarthritis.

Cats : For the control of postoperative pain and inflammation associated with orthopedic surgery, ovariohysterectomy and castration when administered prior to surgery.

Cattle : For use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs in Cattle. For use in diarrhoea in combination with oral re-hydration therapy to reduce clinical signs in Calves of over one week of age and young, non-lactating Cattle. For adjunctive therapy in the treatment of acute mastitis, in combination with antibiotic therapy.

Horses : For use in the alleviation of inflammation and relief of pain in both acute and chronic musculo-skeletal disorders. For the relief of pain associated with equine colic.

DOSAGE AND ADMINISTRATION :

Carefully consider the potential benefits and risk of MELOXIN - I and other treatment options before deciding to use MELOXIN - I. Use the lowest effective dose for the shortest duration consistent with individual response.

Dogs: MELOXIN - I Injection should be administered initially as a single dose at 0.09 mg/lb (0.2 mg/kg) body weight Intravenously (IV) or Subcutaneously (SC), followed, after 24 hours.

Cats: Administer a single, one-time Subcutaneous dose of MELOXIN - I Injection to Cats at a dose of 0.14 mg/lb (0.3 mg/kg) body weight. Use of additional Meloxicam Injection or other NSAIDs is contraindicated. To ensure accuracy of dosing, the use of a 1 ml graduated syringe is recommended.

Cattle: Single Subcutaneous or Intravenous Injection at a dosage of 0.5 mg Meloxicam/kg body weight (i.e. 10 ml/100 kg body weight) in combination with antibiotic therapy or with oral re-hydration therapy, as appropriate. The recommended maximum volume to be administered at a single Injection site is 10 ml.

Horses: Single Intravenous Injection as a dosage of 0.6 mg Meloxicam/kg body weight (i.e. 12 ml/100 kg body weight). For use in the alleviation of inflammation and the relief of pain in both acute and chronic musculo-skeletal disorders.

WITHDRAWAL PERIOD :

Cattle : Meat and Offals: 15 days, Milk : 5 days

Horses & Swine : Meat and Offals : 5 days

PACK SIZE :

30 ml, 100 ml, 250 ml Vial.

Manufactured in India by :

IPPL
Intracin
Pharmaceuticals Pvt. Ltd.
C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

