



PAVEX PLUS

PARVAQUONE AND FUROSEMIDE INJECTION

COMPOSITION :

Each ml contains :

Parvaquone		150 mg
Furosemide	B.P.	55 mg
Sorbitan mono oleate	B.P.	100 mg
(As preservative)		
Oily base		q.s.

INDICATIONS :

A potent injection treatment for theileriosis (East Coast Fever, ECF), particularly for advanced cases with pulmonary oedema. Parvaquone is theilericidal, acting on schizonts and piroplasms; Furosemide is a diuretic, which resolves pulmonary oedema.

DOSAGE AND ADMINISTRATION :

Inject by the Intramuscular route into the neck muscles at the rate of 1 ml/15 kg (10.0 mg Parvaquone, 3.6 mg Furosemide per kg). Repeat after 48 hours. In cases of exceptionally severe infections with pulmonary oedema, further treatment at half the dosage rate may be required at 24 hour intervals.

Calves : 1 ml per 15 kg bodyweight.

Young-stock and adults :

Bodyweight	Dose
100 kg	7 ml
150 kg	10 ml
200 kg	14 ml
250 kg	17 ml
300 kg	20 ml

Alternatively, when pulmonary symptoms have been resolved by Pavex Plus, treatment to achieve parasitic cure may be completed with Pavex, which contains only Parvaquone, at a dose rate of 1 ml/15 kg every 48 hours. Do not use by the Intravenous or Subcutaneous routes. Normal aseptic precautions should be observed.

WITHDRAWAL PERIOD :

Milk : 14 days, Meat : 28 days

PACK SIZE :

100 ml Vial

Manufactured in India by :



Intracin

Pharmaceuticals Pvt. Ltd.

C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

