



TROXIN 3 G

CEFTRIAZONE SODIUM FOR INJECTION B.P.



COMPOSITION :

Each vial contains:

Combipack of A & B contains :

- | | | |
|--------------------------------|------|-------|
| A. Ceftriazone Sodium | B.P. | |
| equi. to Ceftriazone anhydrous | | 3 g |
| B. One ampoule containing | | |
| Sterile Water for Injection | I.P. | 10 ml |

INSTRUCTION FOR USE :

Prepare solution by dissolving the content of vial in 9.6 ml of sterile Water for Injection I.P.

INDICATIONS :

For the treatment of mastitis, respiratory tract infection, skin and soft tissue infection.

DOSAGE AND ADMINISTRATION :

I.M. / I.V. USE

TROXIN 3g Injection should be diluted with 10 ml SWFI and injected at 12-24 hours interval.

It can be administered either Intramuscularly or Intravenously.

Large animals : 7.5 - 25 ml/50 kg bodyweight.

Calves and Goats : 1.5 - 5 ml/10 kg bodyweight.

Dogs : 1.5 - 5 ml/10 kg bodyweight.

Cats : 0.25 - 0.5 ml/kg bodyweight.

WITHDRAWAL PERIOD :

Milk : 7 days

Meat : 28 days

PACK SIZE :

3 gram in 20 ml Vial.

Manufactured in India by :

IPPL
Intracin
Pharmaceuticals Pvt. Ltd.
C-1, B-52,53,56, G.I.D.C. ESTATE,
NADIAD - 387 001, (GUJARAT) INDIA.

