AT WS

Amoxycillin Trihydrate Powder for oral administration

COMPOSITION:

Each gram contains:

Amoxycillin trihydrate 200 mg Excipients ad 1 g

DESCRIPTION:

Amoxycillin is a semisynthetic broad-spectrum penicillin with a bactericidal action against both Gram-positive and Gram-negative bacteria. The spectrum of amoxycillin includes Campylobacter, Clostridium, Corynebacterium, E. coli, Erysipelothrix, Haemophilus, Pasteurella, Salmonella, penicillinase-negative Staphylococcus and Streptococcus spp. The bactericidal action is due to inhibition of cell wall synthesis. Amoxycillin is mainly excreted in urine. A major part can also be excreted in bile.

INDICATIONS:

Gastrointestinal, respiratory and urinary tract infections caused by amoxycillin sensitive micro-organisms, like Campylobacter, Clostridium, Corynebacterium, E. coli, Erysipelothrix, Haemophilus, Pasteurella, Salmonella, penicillinase-negative Staphylococcus and Streptococcus spp. in Calves, Goats, Poultry, Sheep and Swine.

CONTRA-INDICATIONS:

Hypersensitivity to amoxycillin.

Administration to animals with a seriously impaired renal function.

Concurrent administration of tetracyclines, chloramphenicol, macrolides and lincosamides.

Administration to animals with an active microbial digestion.

SIDE EFFECTS:

Hypersensitivity reactions may occur.

DOSAGE AND ADMINISTRATION:

For oral administration:

Calves, Goats and Sheep: Twice daily, 5g / 100 kg body weight for 3-5 days.

Poultry and Swine: 1 kg per 1000-2000 litres of drinking water for 3-5 days.

Note: for pre-ruminant Calves, Lambs and Kids only.

WITHDRAWAL PERIOD:

For meat:

Calves, Goats, Sheep and Swine: 8 days. Poultry: 3 days.



Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE, NADIAD-387001 (GUJARAT) INDIA. www.intracin.com

