

BIPROCIN

Probiotic Feed Additive

COMPOSITION :

Each gram contains 9 different species of beneficial microflora each of 4×10^8 CFU

Bifidobacterium bifidum
Lactobacillus acidophilus
Lactobacillus bulgaricus
Lactobacillus casei
Lactobacillus plantarum
Streptococcus faecium
Saccaromyces cerevisiae
Yeasts : torulopsis spp.
 : aspergillus oryzae
Fortified with phytase & spirulina

INDICATIONS :

- Triggers development of intestinal mucous glands and villi in large numbers and size.
- Minimizes different kinds of stress such as debeaking, vaccination and summer stress.
- Helps to maintain healthy gastrointestinal after antibiotic therapy.
- Reduces the incidents of chick mortality.
- Quicker detoxification of mycotoxins.

DOSAGE AND ADMINISTRATION :

Chicks, Growers & Broilers : 500 g/tonne of feed.

Layers & Breeders : 1 kg/tonne of feed.

Aqua Culture species : 1.5 kg/tonne of feed or as directed by the veterinarian.


Intracin
Pharmaceutical Pvt. Ltd.
C-1,B-53,G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com

