

# BSTAC WS

## Anticoccidial Powder for oral administration

### COMPOSITION :

#### Each 100 gram contains :

Amprolium Hydrochloride	17.5 g
Sulfaquinoxaline	17.5 g
Vitamin A1 retinol acetate	1,500,000 I.U
Vitamin K3	500 mg
Excipients	q.s.

### INDICATIONS :

Coccidiosis in Calves, Sheep, Goat and Poultry

### DOSAGE AND ADMINISTRATION :

For oral administration :

Calves, Sheep and Goats

Preventive : 1 g per 50-100 kg body weight through drinking water or milk for 21 days.

Poultry : 20 g per 20-40 litres of drinking water for 5-7 days

Curative : 5 g per 25-50 kg body weight through drinking water or milk for 5 days.

### WITHDRAWAL PERIOD :

For meat :

Calves, Sheep and Goats : 14 days.

Poultry : 14 days.

### STORAGE :

Store at a temperature not exceeding 25°C.

Protect from light.

### NOTE :

For peruminant Calves, Lambs and only do not administer to hens producing eggs for humans consumption.



## Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,  
NADIAD-387001 (GUJARAT) INDIA.  
[www.intracin.com](http://www.intracin.com)

