

# CLOSIN 10%

## Closantel 10% w/v Injection



### COMPOSITION :

#### Each ml contains :

Closantel	100 mg
Benzyl alcohol B.P	4 % v/v
Propylene glycol B.P	q.s.

### INDICATION :

CLOSIN 10 % is used for the control and treatment of internal parasites including adult and juvenile liver flukes (6-8 weeks), and external parasites. CLOSIN 10 % is indicated for the effective treatment and control of the following parasites - Gastrointestinal roundworms (adult and fourth-stage larvae), *Ostertagia ostertagi*, *O. lyrata*, *Haemonchus placei*, *Trichostongylus axei*, *T. colubriformis*, *Cooperia oncophora*, *C. punctata*, *C. pectinata*, *Bunostomum phlebotomum*, *Nematodirus helvetianus* (adult only), *Napathiger* (adult only), *Oesophagostomum radiatum*.  
Lungworms (adult and fourth-stage larvae): *Dictyocaulus viviparus*.  
Liver flukes (adult and juvenile) (6-8 weeks).  
Cattle grub (parasitic stages): *Hypoderma bovis*, *H. lineatum*.  
Sucking Lice: *Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*.  
Mange Mites (Cattle scab): *Psoroptes ovis*, *Sarcoptes scabiei* var. *bovis*.

### DOSAGE AND ADMINISTRATION :

Target Species :

Cattle , Sheep :

1 ml / 50kg of bodyweight by Subcutaneous Injection only

### WITHDRAWAL PERIOD :

Meat : 49 days

### STORAGE :

Store at a temperature not exceeding 30°C.

Protect from sunlight.

IPPL

## Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,  
NADIAD-387001 (GUJARAT) INDIA.  
[www.intracin.com](http://www.intracin.com)

