

CLOSIN 5%

Closantel 5% w/v Injection

COMPOSITION :

Each ml contains :

Closantel	50 mg
Benzyl alcohol B.P	2% v/v
Propylene glycol B.P	q.s.

INDICATION :

CLOSIN is used for the control and treatment of internal parasites including adult and juvenile liver flukes (6-8 weeks), and external parasites.

CLOSIN is indicated for the effective treatment and control of the following parasites - Gastrointestinal roundworms (adult and fourth-stage larvae), *Ostertagia ostertagi*, *O. lyrata*, *Haemonchus placei*, *Trichostongylus axei*, *T. colubriformis*, *Cooperia oncophora*, *C. punctata*, *C. pectinata*, *Bunostomum phlebotomum*, *Nematodirus helvetianus* (adult only), *Napathiger* (adult only), *Oesophagostomum radiatum*.

Lungworms (adult and fourth-stage larvae): *Dictyocaulus viviparus*.

Liver flukes (adult and juvenile) (6-8 weeks).

Cattle grub (parasitic stages): *Hypoderma bovis*, *H. lineatum*.

Sucking Lice: *Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*.

Mange Mites (Cattle scab): *Psoroptes ovis*, *Sarcoptes scabiei* var. *bovis*.

DOSAGE AND ADMINISTRATION :

Target Species :

Cattle, Sheep : Bodyweight of 50 kg

Dosage : 2-5 ml

WITHDRAWAL PERIOD :

Meat : 49 days

STORAGE :

Store at a temperature not exceeding 30°C.

Protect from sunlight.



Intracin

Pharmaceutical Pvt. Ltd.

C-1, B-53, G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.

www.intracin.com

