

DIMINACIN R.T.U.

Diminazene Aceturate 7% Injection



COMPOSITION :

Each ml contains :

Diminazene Aceturate	70 mg
Phenazone B.P (ANTIPYRINE)	375 mg
Water for Injection B.P	q.s.

INDICATIONS :

Tryptansomiasis due to Trypanosoma congolense, T. Vivax and Brucei, Pyroplasmosis due to Babesia bovis, B. bigemina, B. ovis, B. motasi, B. canis, Theileriosis due to theileria annulata.

CONTRA-INDICATIONS :

Hypersensitivity to diminazene or phenazone.
Administration to animals with an impaired renal and/or hepatic function.

DOSAGE AND ADMINISTRATION :

The Recommended dosage is 3.5 mg of Diminazene aceturate per kg body weight as a single Injection which is 5 ml per 100 kg body weight. Do not exceed a total dose of 4 g of active ingredient in any animal (or 56 ml of DIMINACIN).

ADMINISTRATION :

Deep Intramuscular Injection.

WITHDRAWAL PERIOD :

Meat : 21 days
Milk : 3 days

STORAGE :

Store at a temperature not exceeding 30°C.
Protect from light.



Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com

