

DORACIN

Doramectin Injection 1% w/v

COMPOSITION :

Each ml contains :

Doramectin	10 mg
Oil base	q.s.

INDICATIONS :

Cattle: For treatment and control of gastrointestinal nematodes, lungworms, eyeworms, warbles, lice, mange mites and ticks. The product may also be used as an aid in the control of *Nematodirus helvetianus*, biting lice (*Damalinea bovis*), the tick *Ixodes ricinus* and the mange mite *Chorioptes bovis*.

Sheep: For treatment and control of gastrointestinal roundworms, mange mites and nasal bots.

Pig: For treatment of mange mites, gastrointestinal roundworms, lungworms, kidney worms and sucking lice in Pigs.

DOSAGE AND ADMINISTRATION :

For the treatment and control of gastrointestinal roundworms, lungworms, eyeworms, warbles, lice and mange mites in Cattle, and gastrointestinal roundworms and nasal bots in Sheep, a single treatment of 1 ml (10 mg Doramectin) per 50 kg bodyweight, equivalent to 200 mcg/kg bodyweight, administered in the region of the neck by subcutaneous injection in Cattle and by intramuscular injection in Sheep. For the treatment of clinical signs of *Psoroptes ovis* (Sheep Scab) and elimination of living mites on Sheep a single treatment of 1 ml per 33 kg bodyweight, equivalent to 300 mcg/kg bodyweight, administered in the neck by intramuscular injection. In addition, adequate bio-security measures should be implemented to prevent reinfestation. It is important to ensure that all Sheep which have been in contact with infested Sheep are treated. For the treatment of *Sarcoptes scabiei* and gastrointestinal nematodes, lungworms, kidney worms and sucking lice in Pig, a single treatment of 1 ml per 33 kg bodyweight, equivalent to 300 mcg/kg bodyweight, administered by intramuscular injection. Piglets weighing 16 kg or less should be dosed in accordance with the following table:

Body weight (kg)	Dose (ml)
Less than 4 kg	0.1 ml
5 - 7 kg	0.2 ml
8 - 10 kg	0.3 ml
11 - 13 kg	0.4 ml
14 - 16 kg	0.5 ml

To ensure administration of a correct dose, bodyweight should be determined as accurately as possible; accuracy of the dosing device should be checked. If animals are to be treated collectively rather than individually, they should be grouped according to their bodyweight and dosed accordingly, in order to avoid under- and over- dosing. Maximum injection volume for each target species:

Cattle: 5 ml per injection site.

Sheep: 1.5 ml per injection site.

Pig: 2.5 ml per injection site

CONTRA-INDICATIONS :

Do not use in Dogs, as severe adverse reactions may occur. In common with other avermectins, certain breeds of Dog, such as collies, are especially sensitive to Doramectin and particular care should be taken to avoid accidental consumption of the product. Do not use in case of hypersensitivity to the active substance or any of the excipients.

WITHDRAWAL PERIOD :

Cattle: Meat and offal: 70 days

Not permitted for use in lactating animals producing milk for human consumption. Do not use in pregnant Cows or Heifers, which are intended to produce milk for human consumption, within 2 months of expected parturition.

Sheep: Meat and offal: 70 days

Not permitted for use in lactating animals producing milk for human consumption. Do not use in pregnant Ewes, which are intended to produce milk for human consumption, within 70 days of expected parturition.

Pig: Meat and offal: 77 days

STORAGE :

Store at a temperature not exceeding 30°C.

Protect from light.

IPPL

Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,

NADIAD-387001 (GUJARAT) INDIA.

www.intracin.com