

FLASIN 20%

Tylosin Injection 20% w/v



COMPOSITION :

Each ml contains :

Tylosin Base B.P	200 mg
Benzyl alcohol B.P	2% v/v
Propylene Glycol B.P and	
Water for Injection B.P (1:1)	q.s.

INDICATIONS :

Cattle : Tylosin Injection is indicated for use in the treatment of bovine respiratory complex (shipping fever, pneumonia) usually associated with *Pasteurella multocida* and *Actinomyces pyogenes*; foot rot (necrotic pododermatitis) and calf diphtheria caused by *Fusobacterium necrophorum* and metritis caused by *Actinomyces pyogenes*.

Sheep : Pneumonia, arthritis, contagious and acute agalaxia due *Mycoplasma agalactiae*, mastitis.

Goat : Contagious pleuropneumonia due *Mycoplasma mycoides*.

CONTRA-INDICATIONS :

Not to be used in animals known to be hypersensitive to the active ingredient.

DOSAGE AND ADMINISTRATION :

Cattle:

Inject 5-10 mg/kg of body weight daily intramuscularly.
Do not inject more than 10 ml per injection site.

Sheep and Goats:

Inject 10 mg/kg of body weight daily intramuscularly.
Do not inject more than 5 ml per injection site.

WITHDRAWAL PERIOD :

Cattle :

Edible tissue : 7 days
Milk : 72 hours

Sheep, Goats :

Edible tissue : 7 days
Milk : 48 hours

STORAGE :

Store at a temperature not exceeding 30°C.
Protect from light.



Intracin

Pharmaceutical Pvt. Ltd.

C-1, B-53, G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com

