



# I-TIN PLUS

## Ivermectin and Clorsulon Injection

### COMPOSITION :

Each ml contains :

- Ivermectin B.P. 10 mg
- Clorsulon U.S.P. 100 mg
- Benzyl Alcohol B.P. 1.5% w/v
- Propylene Glycol B.P. q.s.

### INDICATION :

For the treatment and control of : Gastrointestinal immature and adult roundworms.

Lung Worms : Both adult and Fourth - Stage larvae of Dictyocaulus viviparous.

Liver Flukes : Immature and adult Fasciola hepatica & Fasciola gigantica.

Eye Worm : Thelazia spp. Ectoparasites (Dermatobia hominis, Psoroptes bovis) Including Sucking Lice, Ticks & Mites.

### CONTRAINDICATION :

Administration to lactating animals and pregnant heifers within 60 days of Calving. This product is not for intravenous or intramuscular use.

### DOSAGE AND ADMINISTRATION :

For subcutaneous administration.

General : 1.0 ml per 50 kg bodyweight

Cattle : 1 ml per 50 kg bodyweight

Sheep & Goat : 0.5 ml per 25 kg bodyweight

### WITHDRAWAL PERIOD :

For meat : 35 days

### STORAGE :

Store at a temperature not exceeding 30°C.

Protected from light.



**Intracin**

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,  
NADIAD-387001 (GUJARAT) INDIA.  
[www.intracin.com](http://www.intracin.com)

