

I-TIN

Ivermectin Injection 1% B.P

COMPOSITION :**Each ml contains :**

Ivermectin B.P	10 mg
Benzyl Alcohol	
(as preservative) B.P	1.5% v/v
Propylene Glycol B.P	q.s.

INDICATIONS :

For treatment and control of Gastrointestinal nematodes, Lung worms, Eye worms, Warbies and Ectoparasites like mites, ticks and lice in Cattle, Buffaloes, Horses, Sheep, Goats, Camels and Dogs.

CONTRA-INDICATIONS :

I-TIN should not be used Intravenously or Intramuscularly. Do not use in lactating Cows or in dairy Cow during 28 days prior to calving.

DOSAGE AND ADMINISTRATION :

Cattle : 1 ml per 50 kg bodyweight (based on a recommended level of 200 mcg Ivermectin per kg bodyweight).

Administration : Inject Subcutaneously in front or behind the shoulder.

Swine : Dosage : 1 ml per 33 kg bodyweight (based on a recommended level of 303 mcg Ivermectin per kg bodyweight).

For the treatment of Psoroptes Ovis (mange mite), two Injections with a seven day interval are required.

Administration :

Inject Subcutaneously into the neck a sterile 17 gauge 1/2 inch needle in recommended.

WITHDRAWAL PERIOD :

Cattle : Must not be treated within 21 days of slaughter for human consumption. Do not use in Cattle producing milk for human consumption or in dairy cows within 28 days prior to calving.

Swine : Must not be treated within 28 days of slaughter for human consumption.

Sheep : Must not be treated within 21 days of slaughter for human consumption, do not treat lactating Cows or within 21 days before lambing where milk is to be used for human consumption.

STORAGE :

Store at a temperature not exceeding 30°C.
Protect from light.



Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com

