

IMOLY 30% LA

Oxytetracycline Injection

Long Acting 30% w/v



Veterinary Medicines - Injectables

COMPOSITION :

Each ml contains :

Oxytetracycline Dihydrate B.P
eq. to Anhydrous Oxytetracycline 300 mg
in 2-Pyrrolidone Vehicle system q.s.

INDICATIONS :

IMOLY 30% is indicated in the treatment and control of disease in Cattle, Sheep and Pigs caused by, or associated with, organisms sensitive to Oxytetracycline.

Cattle : For the treatment and control of pasteurellosis and pneumonia caused by Oxytetracycline sensitive organisms and as an aid in the treatment of infectious bovine keratoconjunctivitis (New Forest Disease) due to Oxytetracycline sensitive strains of *Moraxella bovis*. This product may also be of value for foul-in-the-foot.

Pigs : For the treatment of pneumonia caused by *Pasteurella*.

Sheep : For the control of enzootic abortion and pneumonia caused by Oxytetracycline sensitive organisms. This product may be used as an aid in the treatment of foot rot, acute, severe mastitis and infectious ovine keratoconjunctivitis (pink-eye).

CONTRA-INDICATIONS :

Not recommended for Cats, Dogs, Horses and Donkeys.

The use of this product during the period of tooth development including late pregnancy may lead to tooth discoloration.

DOSAGE AND ADMINISTRATION :

I.M./Subcutaneous use

Deep intramuscular use only in large & small animals and subcutaneous in poultry.

In large & small animal & poultry.

1ml / 10kg body wt.

0.25ml / 1kg body wt.

WITHDRAWAL PERIOD :

For meat (standard dose):

Cattle and sheep : 28 days.

Swine : 14 days.

For meat (high dose):

Cattle : 35 days.

Sheep and swine : 28 days.

For milk:

Cattle : 10 days.

Sheep : 8 days.

STORAGE :

Store at a temperature not exceeding 30°C.

Protect from light.


Intracrin
Pharmaceutical Pvt. Ltd.

C-1, B-53, G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracrin.com

