



# IMOLY

## Oxytetracycline Spray

### COMPOSITION :

#### Each container contains :

Oxytetracycline hydrochloride 5 g

### INDICATIONS :

IMOLY Spray indicated for the treatment and control of Foot rot, digital dermatitis, Udder's teat, Lips and all skin infections, skin burns, post surgical infections and excoriations caused by Oxytetracycline : sensitive organisms in Cattle, Sheep and animal species.

### DOSAGE AND ADMINISTRATION :

IMOLY Spray suspension for topical use. Shake well before use. Remove any dead tissue dried secretions and debris from the infected site. Spray the product onto the affected area for 1-2 seconds, at a distance of 15-20cm, until an even distribution of the dye is obtained. Repeat the treatment every 12 hours, until complete recovery. Wash spray button with water in case of blockage.

### PRECAUTIONS :

- Pressurized container, protect from sunlight and do not expose to temperatures exceeding 50°C, keep away from sources of ignition, No smoking.
- Do not spray on a naked flame or any incandescent materials.
- Do not pierce or burn even when empty.
- Do not spray on food, food utensils, toys, etc.
- Keep out of reach of children
- Use only as directed.
- Do not use empty can for any purpose; instead dispose it in an environmentally friendly manner.
- Extremely flammable.

### WARNINGS :

- For external use only.
- Use only in a well ventilated area.
- Do not spray in or near the eyes.
- Operators should wear impervious rubber gloves.
- Wash any splashes immediately.
- Wash hand after use.
- People with known hypersensitivity to Oxytetracycline should avoid contact with this product.

### WITHDRAWAL PERIOD :

Meat : Nil

Milk : Nil

### STORAGE :

Store below 25°C.

IPPL

## Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,  
NADIAD-387001 (GUJARAT) INDIA.  
www.intracin.com

