

INCEFT

Ceftiofur Sodium Powder for Injection 1g/4g (Vet)

COMPOSITION :

Each vial contains :

Ceftiofur Sodium eq. to Ceftiofur 1 g

For Intramuscular Injection in Cattle, Buffaloes, Sheep & Goat. It can be used in lactating dairy animals.

INDICATION :

- Genital infections of bovine (acute metritis, cervicitis, prolapsed related to retention of placenta cases) associated with Arcanobacterium pyogenes, Fusobacterium necrophorum and bacteroids spp.
- Respiratory diseases of Cattle, Buffalo, Sheep and Goat (Shipping fever, pneumonia) associated with Pasteurella haemolytica, Pasteurella multocida & Haemophilus somnus.
- Acute interdigital necrobacillosis (foot rot, pododermatitis) caused by Fusobacterium and Bacteroides.

DOSAGE AND ADMINISTRATION :

Cattle, Sheep and Goat : 1.1 to 2.2 mg/kg b.wt. or 1 to 2 ml (after reconstitution) /45.45 kg b.wt. by IM use only, once daily for 3-5 days.

Buffalo : 2 to 2.4 mg/kg b.wt. by IM use only, once daily for 3-5 days.

Dog : 2.2 mg/kg b.wt. by SC use only, once daily for 5 to 14 days.

Camel : 2.2 mg/kg b.wt. by IM once daily for 3-5 days.

Horse : 2.2 to 4.4 mg/kg b.wt. or 1 to 2 ml (after reconstitution) /22.72 kg b.wt. by IM use only, once daily for 3 to 5 days.

Swine : 3 to 5 mg/kg b.wt. or 1 to 1.7 ml (after reconstitution) / 16.66 kg b.wt. by IM use only, once daily for 3 to 5 days.

Chicken (day-old chick) : 0.08 to 0.2 mg/Chick, SC use only in the neck region (1 g vial after reconstitution with 20 ml water for injection will treat approximately 5,000 to 12,500 day-old chicks)

WITHDRAWAL PERIOD :

For Cattle, Buffalo & Swine :

Milk : Nil | Meat : 4 days

For Sheep & Goat :

Milk : Nil | Meat: Nil

STORAGE :

Store at a temperature not exceeding 30°C.
Protect from light.

IPPL

Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com

