



IROFLOX 100

Enrofloxacin 20% Oral Liquid

COMPOSITION :

Each ml contains :

Enrofloxacin BP Vet 200 mg

INDICATIONS :

Enrofloxacin 20% Oral Solution has a broad spectrum efficacy against a wide range of bacteria and mycoplasma.

In Poultry indicated for :

CRD, Fowl cholera (avian H.S), Infectious coryza, salmonellosis, clostridiosis (necrotic enteritis), colibacillosis, omphilitis and the infections caused by enrofloxacin sensitive bacteria.

DOSAGE AND ADMINISTRATION :

1ml Enrofloxacin 20% Oral Solution per 4 litres of drinking water for 3 days continuously.

In case of salmonella and mycoplasma continue treatment for 5 days. (10 mg Enrofloxacin per kg body weight).

PRECAUTION :

Do not use in case of resistance and sensitivity to quinolones, do not use with NSAIDS, macrolides, chlorempenicols and tetracyclines because of antagonistic interactions of the drugs.

- Consult the veterinarian before use.
- Keep out of the reach of children.

WITHDRAWAL PERIOD :

Meat : 12 days.

STORAGE :

Store below 25°C in a cool and dry place.
Protect from light and moisture.



Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com

