

MARBACIN 10%

Marbofloxacin Injection 10% w/v

COMPOSITION :

Each ml Contains :

Marbofloxacin	B.P.	100 mg
m cresol	B.P.	2 mg
(As Preservatives)		
Water for injection	B.P.	q.s.

INDICATIONS :

Cattle : Treatment of respiratory infections caused by sensitive strains of *Pasteurella multocida*, *Mannheimia (Pasteurella) haemolytica* and *Mycoplasma bovis*.
Treatment of acute mastitis caused by *E. coli* strains sensitive to marbofloxacin during the lactation period.

Sows : Treatment of Metritis Mastitis Agalactia Syndrome caused by bacterial strains sensitive to marbofloxacin.

CONTRA-INDICATIONS :

Bacterial infections with resistance to other fluoroquinolones (cross resistance).
Do not administer to an animal previously found to be hypersensitive to marbofloxacin or other quinolone.

DOSAGE :

Cattle : Treatment of respiratory infections: The recommended dosage is 2 mg/kg (1 ml/50kg bw) in a single daily injection by subcutaneous or intramuscular route, for 3 to 5 days.

Treatment of acute mastitis : The recommended dosage is 2 mg/kg (1 ml/50kg bw) in a single daily injection by subcutaneous or intramuscular route, for 3 consecutive days. The first injection may also be given by the intravenous route.

Sows : The recommended dosage is 2 mg/kg (1 ml/50kg bw) in a single daily injection by the intramuscular route, for 3 days.

Calves : It has been used to treat respiratory infections (bovine respiratory disease) at 2 mg/kg for 3–5 days or a single dose administered once at 8–10 mg/kg.

WITHDRAWAL PERIOD :

Cattle :

Meat and offals : 6 days

Milk : 36 hours

Sows :

Meat and offals : 4 days

STORAGE :

Store at a temperature not exceeding 30°C.

Protect from light.



Intracin

Pharmaceutical Pvt. Ltd.

C-1, B-53, G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com