NEROCIN

Injection of Methylcobalamin, Vitamin B6 & Nicotinamide

COMPOSITION: Each ml contains:

Methylcobalamin 500 mcg Pyridoxine Hydrochloride B.P 50 mg Nicotinamide B.P 50 mg Benzyl Alcohol B.P 2% w/v

(as preservative)

Water for Injection B.P q.s.

Overanges of vitamins are added to compensate loss on storage.

INDICATION:

Liver disorders like fatty liver degeneration and hepatitis. Neurological disorders e.g. Nutritional Neuritis, dermatitis, macrocytic anemia. Debility and exhaustion.

Anorexia, Fatigue.

To prevent anemia and normal RBC production.

As a supportive therapy with antimicrobials and antiparasitic treatment.

DOSAGE AND ADMINISTRATION:

Cattle, Buffalo, Horse : 5 to 10 ml IM or IV daily Dog : 2 to 3 ml IM or IV daily

For normal Condition: 3 days For Acute Condition: 7 days

WITHDRAWAL PERIOD:

Milk : Nil Meat : Nil

STORAGE:

Store at a temperature not exceeding 30°C.

Protect from light.



Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE, NADIAD-387001 (GUJARAT) INDIA. www.intracin.com

