# **PAXON PLUS**

## **Parvaquone And Furosemide Injection**

**COMPOSITION:** 

Each ml contains:

Parvaquone 150 mg
Furosemide B.P 55 mg
Sorbitan Mono Oleate B.P 100 mg

(as preservative)

Oil base q.s.

#### **INDICATION:**

PAXON PLUS is a potent Injection treatment for theileriosis (East Coast Fever) particularly for advanced cases with pulmonary oedema. Parvaquone is theilericidal, acting on schizonts and piroplasms: Frusemide is a diuretic, which resolves pulmonary oedema.

#### **DOSAGE AND ADMINISTRATION:**

Inject by the Intramuscular route into the neck muscles at the rate of 1 ml/50 kg (10.0 mg Parvaquone, 3.6 mg Furosemide per kg). Repeat after 48 hours. In case of exceptionally server infections with pulmonary oedema, further treatment at half the dosage rate may be required at 24 hour intervals. Calves: 1 ml/50 kg body weight Young-stock and adults:

Body weight	Dose
100kg	7ml
150kg	10ml
200kg	14ml
250kg	17ml
300kg	20ml

Alternatively. When pulmonary symptoms have been resolved by PAXON PLUS, treatment to achieve parasitic cure may be completed with PAXON which contains only Parvaquone, at a dose rate of 1 ml/50 kg every 48 hours. Do not use by the Intravenous or Subcutaneous route. Normal asceptic precautions should be observed

### WITHDRAWAL PERIOD:

Milk : 14 days Meat : 28 days

#### STORAGE:

Store at a temperature not exceeding 30°C. Protect from light.



## Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE, NADIAD-387001 (GUJARAT) INDIA. www.intracin.com

