



# TRISULFACIN WS

## Sulphadimidine + Sulphadiazine + Sulphathiazole Sodium with Vitamin K3 Powder for Oral Administration

### COMPOSITION :

#### Each gram contains :

Sulphadimidine sodium B.P	125 mg
Sulphadiazine sodium B.P	125 mg
Sulphathiazole sodium B.P	125 mg
Vitamin K3 B.P	10 mg

### INDICATIONS :

It is used for treatment of Gastro-Intestinal diarrhoea, Respiratory tract infections, Shipping fever in large animals and intestinal coccidiosis in Poultry.

### DOSAGE AND ADMINISTRATION :

#### Poultry :

1 g Trisulfacin ws per 2-4 litres of drinking water to be given as follows :  
3 days treatment, 2 days without treatment, then 2 days treatment.

#### Calves, Lambs and Kids :

1 g Trisulfacin ws per 5 kg body weight daily for 3-4 days.

**Warning :** Not to be used in poultry producing eggs for human consumption.

### WITHDRAWAL PERIOD :

Poultry : 5 days

Meat : 21 days

### WARNING :

Not to be used in poultry producing eggs on human consumption.

Veterinary Medicines - Water Soluble Powder



## Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,  
NADIAD-387001 (GUJARAT) INDIA.  
[www.intracin.com](http://www.intracin.com)

