

TROXIN

Ceftriaxone Sodium for Injection B.P



COMPOSITION :

Each vial contains :

Ceftriaxone Sodium B.P
equi. to Ceftriaxone anhydrous 3 g

INDICATION :

Preoperative prophylaxis, post operative treatment, haemorrhagic septicemia, broncho pneumonia, nephritis, cystitis, endometritis, metritis, pyometra, mastitis, actionbacillosis, bacterial meningitis and septicaemia.

CONTRAINDICATION :

Ceftriaxone is contraindicated in animal with known allergy to cephalosporin class of antibiotics. Careful inquiry should be made to determine whether the animal has had a previous history of hypersensitivity reaction to cephalosporins, penicillins or other drugs. Biliary elimination occurs and hence, Ceftriaxone should be given cautiously to those species of animal with extended large intestines e.g. Horse.

DOSAGE AND ADMINISTRATION :

I.M / I.V USE

TROXIN 3g Injection should be diluted with 10 ml WFL and injected at 12-24 hours interval. It can be administered either intramuscularly or intravenously.

Large animals : 7.5 - 25ml / 50kg body weight.

Calf and Goat : 1.5 - 5ml / 10kg body weight .

Dogs : 1.5 - 5ml / 10kg body weight.

Cat : 0.25 - 0.5 ml/kg body weight.

WITHDRAWAL PERIOD :

Milk : 7 days

Meat : 28 days

STORAGE :

Store at a temperature not exceeding 30°C.

Protect from light.

IPPL

Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com

