

TWIN-PGS

Benzyl Penicillin Powder for Injection B.P

COMPOSITION :

Each pack contains :

1. Each vial contains :

Benzyl Penicillin Sodium B.P

equi. to Benzyl Penicillin

5000000 I.U.(3000 mg)

Buffering Agent : Anhydrous Sodium Citrate USP

2. Sterile Water for Injection I.P

15 ml 1-Ampoule

INDICATION :

TWIN-PGS is used in treatment of mastitis, metritis, actinomycosis, anthrax, black leg, tetanus, haemorrhagic septicaemia in Cattle; Strangles, corynebacterial pneumonia in foals, tetanus in Horses; Erycephalus in Pigs; Tetanus, foot rot, malignant oedema in Sheep and Goat; Rickettsiosis, tetanus and wound infection in Dogs.

CONTRAINDICATION :

Do not use in animals with a known hypersensitivity reaction to any of the penicillins.

TWIN-PGS should not be used in Cats, Rabbits, Guinea Pigs, Hamsters, Gerbils or other similar small animals.

DOSAGE AND ADMINISTRATION :

Cats and Dogs : Intramuscular 20,000 to 40,000 I.U. per kg of bodyweight every six to eight hours.

Horses : Intramuscular 20,000 I.U. per kg of bodyweight every six to eight hours.

WITHDRAWAL PERIOD :

Calves, Preruminating : Meat : 7 days

Cattle : Meat : 4 days, Milk : 48 hours

Sheep : Meat : 8 days

Pigs : Meat : 6 days

Animal must not be slaughtered and milk may not be used for human consumption during treatment.

STORAGE :

Store at a temperature not exceeding 30°C.

Protect from light.

IPPL

Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,

NADIAD-387001 (GUJARAT) INDIA.

www.intracin.com

